

# COMBINATORICS OF INTERVAL EXCHANGE TRANSFORMATIONS

## LECTURE NOTES FOR DYNAMIQUE EN CORNOUAILLE

SÉBASTIEN FERENCZI

### 1. PRELIMINARIES

#### 1.1. Interval exchange transformations.

**Definition 1.** Let  $r \geq 2$ . Let  $\Lambda_r$  be the set of vectors  $(\lambda_1, \dots, \lambda_r)$  in  $\mathbb{R}^r$  such that  $0 \leq \lambda_i \leq 1$  for all  $i$  and  $\sum_{i=1}^r \lambda_i = 1$ . An  $r$ -interval exchange transformation, or iet for short, is given by a vector  $\lambda \in \Lambda_r$  and a permutation  $\pi$  of  $\{1, 2, \dots, r\}$ . The map  $T_{\lambda, \pi}$  is the piecewise translation defined by partitioning the interval  $X = [0, 1[$  into  $r$  sub-intervals of lengths  $\lambda_1, \lambda_2, \dots, \lambda_r$  and rearranging them according to the permutation  $\pi$ ; or, formally,

$$T_{\lambda, \pi} x = x + \sum_{j < i} \lambda_{\pi j} - \sum_{j < i} \lambda_j$$

when  $x$  is in the interval

$$X_i = \left[ \sum_{j < i} \lambda_j, \sum_{j \leq i} \lambda_j \right[.$$

This is a class of systems introduced by V. Oseledec [20], which have been extensively studied, by three kind of methods: by definition, they are *one-dimensional* systems, and will be studied as such in Sections 2, 4, 6 below; but the strongest results on iet have been obtained by lifting the transformation to *higher dimensions* and using deep geometric methods; however, most of these results have been reproved by using *zero-dimensional* methods, which we show in Sections 3 and 5, after giving the necessary definitions of word combinatorics and symbolic dynamics in the next two subsections; then Section 7 is devoted to the last big open question on iet. For readers who prefer geometric methods, they will find them in the two excellent courses [27] and [28].

Throughout this course, we denote by  $\beta_i$ ,  $1 \leq i \leq r - 1$ , the  $i$ -th discontinuity of  $T^{-1}$ , namely  $\beta_i = \sum_{j \leq i} \lambda_{\pi j}$ , while  $\gamma_i$  is the  $i$ -th discontinuity of  $T$ , namely  $\gamma_i = \sum_{j \leq i} \lambda_j$ . We shall use also  $\gamma_0 = 0$ ,  $\gamma_r = 1$ . Then  $X_i$  is the interval  $[\gamma_{i-1}, \gamma_i[$ .

**Warning:** roughly half the texts on interval exchange maps re-order the subintervals by  $\pi^{-1}$ ; as it is not always clear to which half a given text belongs, we insist that the present definition corresponds to the following ordering of  $T X_i$ : from left to right,  $T X_{\pi(1)}, \dots, T X_{\pi(r)}$ . It makes sense to re-order also the  $X_i$ , thus defining  $T$  by two permutations  $\pi_0$  and  $\pi_1$  (though of course sometimes  $\pi_0^{-1}$  and  $\pi_1^{-1}$  are used...), see [28] for example.

---

Date: July 4, 2011.

### 1.2. Word combinatorics.

**Definition 2.** We look at finite words on a finite alphabet  $\mathcal{A}$ . A word with  $k$  letters,  $w_1 \dots w_k$ , is of length  $k$ . The concatenation of two words  $w$  and  $w'$  is denoted by  $ww'$ . The empty word is the unique word of length zero.

A word  $w = w_1 \dots w_k$  occurs at place  $i$  in a word  $v = v_1 \dots v_s$  or an infinite sequence  $v = v_1 v_2 \dots$  if  $w_1 = v_i, \dots, w_k = v_{i+k-1}$ . We say that  $w$  is a factor of  $v$ . When it is finite, we denote by  $N(w, v)$  the number of occurrences of  $w$  in  $v$ .

The empty word is a factor of any  $v$ . Prefixes and suffixes are defined in the usual way.

A language  $L$  is a set of words such if  $w$  is in  $L$ , all its factors are in  $L$ , and  $wb$  is in  $L$  for at least one letter  $b$  of  $\mathcal{A}$ .

A language  $L$  is uniformly recurrent if for each  $w$  in  $L$  there exists  $n$  such that  $w$  occurs in each word of length  $n$  of  $L$ .

The language  $L(u)$  of an infinite sequence is the set of all its finite factors.

**Definition 3.** Let  $L$  be a fixed language. A word  $w$  is right special, resp. left special if there exist at least two different letters  $x$  such that  $xw$ , resp.  $wx$ , is in  $L$ .

The complexity of  $L$  is the function  $p_L$  which to each positive integer  $n$  associates the number of different words of length  $n$  in  $L$ .

The Rauzy graph of length  $n$  of  $L$  is the graph whose vertices are the words of length  $n$  in  $L$ , with an edge  $w \rightarrow w'$  if there exists a word  $v$  of length  $n - 1$  such that  $w = av$ ,  $w' = vb$ , and  $avb \in L$ .

Note that the Rauzy graphs should not be confused with the Rauzy diagrams used in [28] to describe the induction on interval exchanges.

### 1.3. Dynamical systems.

**Definition 4.** The symbolic dynamical system associated to a language  $L$  is the one-sided shift  $S(x_0 x_1 x_2 \dots) = x_1 x_2 \dots$  on the subset  $X_L$  of  $\mathcal{A}^{\mathbb{N}}$  made with the infinite sequences such that for every  $t, s$ ,  $x_t \dots x_{t+s-1}$  is in  $L$ .

For a word  $w = w_1 \dots w_k$  in  $L$ , the cylinder  $[w]$  is the set  $\{x \in X_L; x_0 = w_1, \dots, x_{k-1} = w_k\}$ .

$(X_L, S)$  is minimal if  $L$  is uniformly recurrent.

$(X_L, S)$  is uniquely ergodic if there is one  $S$ -invariant probability measure  $\mu$ ; then the frequency of the word  $w$  is the measure  $\mu[w]$ .

Starting from a (in general, geometric in origin) topological dynamical system  $(X, T)$ , we can get a symbolic dynamical system:

**Definition 5.** For a transformation  $T$  defined on a set  $X$ , partitioned into  $X_1, \dots, X_r$ , and a point  $x$  in  $X$ , its trajectory is the infinite sequence  $(x_n)_{n \in \mathbb{N}}$  defined by  $x_n = i$  if  $T^n x$  falls into  $X_i$ ,  $1 \leq i \leq r$ .

The language  $L(T)$  is the set of all finite factors of its trajectories.

The coding of  $(X, T)$  by the partition  $\{X_1, \dots, X_r\}$  is the symbolic dynamical system  $(X_{L(T)}, S)$ .

The natural coding of an  $r$ -interval exchange is its coding by the partition into the intervals  $X_i$ ,  $1 \leq i \leq r$  defined above.

If the transformation  $T$  is minimal (i.e. every orbit is dense), all its trajectories have the same finite factors, and the language  $L(T)$  is uniformly recurrent; the special words depend on the language and not on the individual trajectories; thus they are defined by any trajectory of  $T$ . If there is no periodic orbit, every word  $w$  is a factor of a bispecial word; hence the bispecial words determine the finite factors of the trajectories, and thus the symbolic dynamical system  $(X_{L(T)}, S)$ .

If  $(X_{L(T)}, S)$  is the natural coding of an iet  $T_{\lambda, \pi}$ , it is not topologically conjugate to  $([0, 1[, T_{\lambda, \pi})$ , but it shares all its properties of minimality and unique ergodicity, and any invariant measure for one of these systems can be carried to the other one.

**Definition 6.** *The induced map of any transformation  $T$  on a set  $Y$  is the map  $y \rightarrow T^{r(y)}y$  where, for  $y \in Y$ ,  $r(y)$  is the smallest  $r \geq 1$  such that  $T^r y$  is in  $Y$  (in all cases considered in this course,  $r(y)$  is finite).*

**Definition 7.** *Let  $(X, T, \mu)$  be a finite measure-preserving dynamical system.*

*A complex number  $\theta$  is an eigenvalue of  $T$  (denoted multiplicatively) if there exists a non-constant  $f$  in  $\mathcal{L}^2(X, \mathbb{C})$  such that  $f \circ T = \theta f$  in  $\mathcal{L}^2(X, \mathbb{C})$ ;  $f$  is then an eigenfunction for the eigenvalue  $\theta$ . We consider only non-constant eigenfunctions, thus  $\theta = 1$  is not an eigenvalue if  $T$  is ergodic.  $T$  is weakly mixing if it has no eigenvalue.*

**Definition 8.** *In  $(X, T, \mu)$ , a (Rokhlin) tower is a collection of disjoint measurable sets called levels  $F, TF, \dots, T^{h-1}F$ .*

## 2. MINIMALITY

The question of *minimality* of iet is an old one, for which we find useful to give a quick reminder. The following fundamental lemma is proved in [16] and [15]; note that it stated as an independent lemma in [14], but with the improved, though unfortunately false, bound  $r + 1$ .

**Lemma 1.** *The induced map  $S_J$  of an  $r$ -iet  $T$  on an interval  $J$  is an  $s$ -iet for some  $s \leq r + 2$ .*

**Proof** We look at the (at most)  $r + 1$  points made by the  $\gamma_i$  and the two endpoints of  $J$ ; if  $x$  is any of these points, let  $s(x)$  be the largest negative integer  $s$  such that  $T^s x$  is in the interior of  $J$ ; we partition  $J$  by the (at most)  $r + 1$  points  $T^{s(x)}x$ . Then on each of these (at most)  $r + 2$  intervals  $S_J$  is continuous and is of the form  $T^j$  for a constant  $j$ .  $\square$

**Definition 9.**  *$T_{\lambda, \pi}$  satisfies the i.d.o.c. property [16] if the orbits of the discontinuity points  $\gamma_i$ ,  $1 \leq i \leq r - 1$ , are infinite and disjoint.*

**Proposition 2.** *The i.d.o.c. condition implies minimality.*

### Proof

We show first that there is no periodic point: if  $T^m x = x$ , let  $b$  be the  $T^n \gamma_j$  nearest to  $x$  on the left, for  $0 \leq j \leq r - 1$ ,  $0 \leq n \leq m - 1$ . Then  $T^m b = b$  as each  $T^i$ ,  $1 \leq i \leq m - 1$ , is an isometry on  $[b, x]$ ; this contradicts the i.d.o.c. unless  $b = T^a 0$ , and  $0$  is itself an image of a discontinuity if  $\pi 1 \neq 1$ , while  $\pi 1 = 1$  contradicts also the i.d.o.c. (in different ways, depending whether  $\pi 2 \neq 2$ , or  $\pi 2 = 2$  and  $\pi 3 \neq 3$ , etc...).

Given an interval  $J$ , we make the *induction castle* of  $J$ :  $J$  is partitioned into  $s$  subintervals  $J_t$ ,  $1 \leq t \leq s$ , the  $T^j J_t$ ,  $1 \leq t \leq s$ ,  $0 \leq j \leq h_t - 1$ , are disjoint intervals,  $T^{h_t} J_t = S J_t \subset J$ . Let  $Y$  be  $\cup_{1 \leq t \leq s} \cup_{0 \leq j \leq h_t - 1} T^j J_t$ ;  $Y$  is a union of intervals, let  $G$  be the union of their left ends. Then  $TY = Y$ , and for  $x \in G$ , either  $Tx \in G$  or  $x = \gamma_j$  for  $0 \leq j \leq r - 1$ . Because  $G$  is finite and  $T$  aperiodic, for all  $x \in G$  there exists  $n$  such that  $T^n x = \gamma_j$  for  $0 \leq j \leq r - 1$ . Similarly for  $x \in G$ , either  $T^{-1}x \in G$  or  $T^{-1}x = \beta_j$  for  $1 \leq j \leq r - 1$ , and there exists  $m$  such that  $T^{-m}x = \beta_j$  for  $1 \leq j \leq r - 1$ . The only possibility for  $x$  which does not contradict the i.d.o.c. is  $x = \gamma_0$ , thus  $Y = X$ .

Now if the orbit of  $x$  is not dense, its closure does not intersect an interval  $J$ , which contradicts the fact that the induction castle of  $J$  fills  $X$ .  $\square$

It is well known and stated in [16] that, if the permutation  $\pi$  is *irreducible* ( $\pi\{0, \dots, l\} \neq \{0, \dots, l\}$  if  $l \neq r$ ), then *total irrationality* (the  $\lambda_i$  satisfy no rational relation except  $\sum \lambda_i = 1$ ) implies the i.d.o.c. condition. But the i.d.o.c. condition is strictly weaker than total irrationality: for three intervals it means that  $\lambda_1$  and  $\lambda_2$  do not satisfy any rational relation of the forms  $p\lambda_1 + q\lambda_2 = p - q$ ,  $p\lambda_1 + q\lambda_2 = p - q + 1$ , or  $p\lambda_1 + q\lambda_2 = p - q - 1$ , for  $p$  and  $q$  integers. Also, the i.d.o.c. condition is not equivalent to minimality, here is a counter-example from [27]:  $k = 4$ ,  $\pi = (4321)$ ,  $\lambda_1 = \lambda_3$ ,  $\lambda_2 = \lambda_4$ ,  $\frac{\lambda_1}{\lambda_2} = \frac{\lambda_3}{\lambda_4}$  is irrational.

The i.d.o.c. condition ensures that  $(X_u, S)$  and  $(X, T_{\lambda, \pi})$  are minimal, each natural coding  $u$  is uniformly recurrent and the language  $L_u$  is the same for all the natural codings.

**Proposition 3.** *The language of the natural coding of an  $r$ -iet satisfying the i.d.o.c. condition has complexity  $(r - 1)n + 1$  for all  $n \geq 0$ .*

**Proof**

The cylinder  $[w_1 \dots w_n]$  is the set  $\cap_{i=0}^n T^{-i} X_{w_{i+1}}$ . By induction on  $n$ , these are either empty sets or intervals, and the partition of  $X$  into nonempty cylinders of length  $n$  is the partition of  $X$  by the points  $T^{-i} \gamma_j$ ,  $1 \leq j \leq r - 1$ ,  $0 \leq i \leq n - 1$ . The i.d.o.c. condition ensures that all these points are different.  $\square$

Note that the left special words of length  $n$  are the prefixes of length  $n$  of the trajectories of the  $\beta_i$ ,  $1 \leq i \leq r - 1$ . Thus when  $n$  is large enough, there are  $r - 1$  left special words of length  $n$ , with two extensions for each one - and the same for right special words by looking at  $T^{-1}$ .

### 3. INVARIANT MEASURES

We shall need the fundamental

**Lemma 4.**  $p_L(n + 1) - p_L(n) = \sum_{w \in RS_n} (\#D(w) - 1)$  where  $RS_n$  is the set of right special words of length  $n$  and  $D(w)$  is the set of letters  $a$  such that  $wa$  is in  $L$ .

and, as a consequence, the result of M. Morse and G. Hedlund [19]

**Proposition 5.** *If  $p_L(n) \leq n$  for at least one  $n$ , then  $L$  is the union of a finite number of  $L(w^j)$  where each infinite sequence  $w^j$  is ultimately periodic, (namely, there exist positive integers  $n_j$  and  $t_j$  such that  $w_{n+t_j}^j = w_n^j$  for all  $n > n_j$ ), and  $p_L(n)$  is bounded.*

**Proof**

Then either  $p(1) = 1$  or there exists  $m$  with  $p(m + 1) = p(m)$ . There is no right special word of length  $m$ , and a loop in each connected component of the Rauzy graph.  $\square$

**Proposition 6.** *A minimal symbolic system such that  $p(n + 1) - p(n) = r$  for all  $n$  has at most  $r - 1$  ergodic invariant measures.*

**Proof**

We begin by showing that there are at most  $r$  invariant measures. Let  $d_{n,1}, \dots, d_{n,r}$  be the right special words of length  $n$ , possibly with repetitions. For a word  $w$ , we define  $\nu_{n,i}(w) = \lim_{u \rightarrow +\infty} \frac{N(w, d_{n,i}^u)}{u |d_{n,i}|}$ . By taking subsequences, we ensure the  $\nu_{n,i}$  converge to a probability  $\mu_i$  on  $X_L$ ,  $1 \leq i \leq r$ .

We remark that if  $n$  is large enough every word in  $L$  of length at least  $(r + 2)n$  contains one of the  $d_{n,i}$ , because such a word contains at least  $(r + 1)n$  factors of length  $n$ , hence two must be equal, and if there is no right special words among them this creates a loop in the Rauzy graph thus  $X_L$

contains ultimately periodic sequences, which is impossible for a minimal system of complexity  $rn + s$ .

Let  $\mu$  be an ergodic invariant probability on  $X_L$ . By the ergodic theorem, we choose an  $x \in X_L$  such that for every  $w \in L$ ,  $\mu([w]) = \lim_{u \rightarrow +\infty} \frac{N(w, x_0 \dots x_{u-1})}{u}$ . We fix  $n$  and cut  $x$  into disjoint words of length  $(r+2)n$ , each of which contains a  $d_{n,i}$ . Thus there exists  $t(n)$  such that  $d_{n,t(n)}$  occurs in the  $j$ -th word for a set of  $j$  of upper density at least  $\frac{1}{r}$ , and we choose  $t$  such that  $t(n) = t$  for infinitely many  $n$ . For those  $n$  and  $m$  large,  $N(w, x_0 \dots x_{m(r+2)n-1}) \geq \frac{m}{2r} N(w, d_{n,t})$ . Dividing by the lengths and letting  $m$  then  $n$  tend to infinity we get  $\mu([w]) \geq \frac{1}{2r(r+2)} \mu_t([w])$ . Thus  $\mu = c\mu_t + (1-c)\mu'$  for some positive measure  $\mu'$  and, as ergodic measures are extremal,  $\mu = \mu_t$ .

To improve the bound, we notice that if for infinitely many  $n$  there are at most  $r-1$  right special factors of length  $n$ , the above reasoning implies that there are at most  $r-1$  ergodic invariant measures. Thus we can assume that for each  $n$  large enough there are exactly  $r$  right special words, and thus by Lemma 4 each of them can be extended by two letters only. We shall show now that there exist  $K$  and  $1 \leq j \leq r$  such that for infinitely many  $n$  every word in  $L$  of length at least  $Kn$  contains one of the  $d_{n,i}$ ,  $i \neq j$ ; then again the above reasoning implies our result.

Indeed, we suppose the above assertion is not satisfied; then for every  $j$ , and  $n$  large enough, there is a path in the Rauzy graph of length  $n$ , going from  $d_{n,j}$  to  $d_{n,j}$ , not meeting any  $d_{n,i}$  except at the two ends; by minimality and because there are only two letters extending  $d_{n,j}$ , this is the only path satisfying these conditions, and our hypothesis implies that it can be followed  $q_{n,j}$  times consecutively, with  $q_{n,j}$  tending to infinity with  $n$ . By following this path, we define a word  $g_{n,j}$  such that  $d_{n,j}g_{n,j}$  begins and ends with  $d_{n,j}$ , with no other occurrence of any  $d_{n,i}$ , and the word  $g_{n,j}^{q_{n,j}}$  occurs in  $L$ ; our hypothesis implies also that  $\frac{q_{n,j}|g_{n,j}|}{n}$  tends to infinity with  $n$ . If we take  $q_{n,j}$  maximal, then  $g_{n,j}^{q_{n,j}-1}$  is right special, thus is identified with some  $d_{n_1,i}$ ; by unicity of the path,  $g_{n_1,i}$  is identified with  $g_{n,j}$ , thus  $q_{n_1,i}$  with  $q_{n,j}$ ; but then  $\frac{q_{n_1,i}|g_{n_1,i}|}{n_1} \leq \frac{q_{n,j}}{q_{n,j}-1}$  is smaller than 2 for some arbitrarily large  $n_1$ , which contradicts our hypothesis.  $\square$

This result is not optimal: the hypothesis can be weakened [3], and the optimal bound for the number of invariant measures for an  $r$ -iet is  $\lfloor \frac{r}{2} \rfloor$  [13][23]. But it is enough for our purpose, as it implies that, under the i.d.o.c. condition, 3-iet are uniquely ergodic, and 4-iet have at most two invariant ergodic measures.

Note that unique ergodicity is a strong notion:

**Proposition 7.** *If  $(X, T)$  is uniquely ergodic,  $\mu$  its invariant probability measure,  $g$  a continuous function on  $X$ , then*

$$\frac{1}{N} \sum_{n=0}^{N-1} g \circ T^n \rightarrow \mu(g) \quad \text{uniformly.}$$

**Proof**

Otherwise, there exist  $\delta > 0$ ,  $f$  continuous, a sequence  $n_k \rightarrow +\infty$  and a sequence of points  $x_{n_k}$  such that

$$\left| \frac{1}{n_k} \sum_{n=0}^{n_k-1} g(T^n x) - \mu(g) \right| > \delta.$$

By compactity, there exists a subsequence  $m_k$  of  $n_k$  such that for every continuous  $g$ ,

$$\lim_{k \rightarrow +\infty} \frac{1}{m_k} \sum_{n=0}^{m_k-1} g \circ T^n$$

exists and defines a measure  $\nu$ .  $\nu$  is then a  $T$ -invariant probability, hence  $\nu = \mu$ , which contradicts the assumption.  $\square$

#### 4. KEANE'S COUNTER-EXAMPLES

It was conjectured by M. Keane [16] that the i.d.o.c. condition implies unique ergodicity for every  $r$ ; when it was made this conjecture had already been disproved by W. Veech [22]. Veech's counter-example was a 5-iet. Then Keane [17] lowered the number of intervals required for a counter-example to four, which is optimal in view of Proposition 6. But his paper uses very different techniques, and there appear for the first time two ideas which were to be named and systematically studied later: one is the *induction*, a different form of which will give the *Rauzy induction* and is the starting point of the geometric methods; the other one is the use of *matrices for adic systems*.

**Lemma 8.** *Let be the 4-iet with vector  $(\lambda_1, \dots, \lambda_4)$  and permutation  $\pi$  sending  $1 \rightarrow 4, 2 \rightarrow 2, 3 \rightarrow 1, 4 \rightarrow 3$  (denoted by (4213)). Suppose*

- $\lambda_1 < \lambda_4 < \lambda_3, \lambda_4 < \lambda_1 + \lambda_2,$
- *for  $1 \leq k < m$   $T^{k-1}[\gamma_1, \beta_1[ \subset X_2$ , then  $T^{m-1}[\gamma_1, \beta_1[ \not\subset X_2$  and  $T^{m-1}[\gamma_1, \beta_1[ \subset X_2 \cup X_3,$*
- *for  $1 \leq k < p$   $T^k[\gamma_2, \gamma_2 + \lambda_4[ \subset X_3$ , then  $T^p[\gamma_2, \gamma_2 + \lambda_4[ \not\subset X_3$  and  $T^p[\gamma_2, \gamma_2 + \lambda_4[ \subset X_3 \cup X_4.$*

*Then the induced map  $S$  of  $T$  on  $[\gamma_3, 1[$  is a 4-iet on an interval of length  $\lambda_4$ , with permutation (2431) and vector  $(\lambda'_4, \lambda'_3, \lambda'_2, \lambda'_1)$  such that, if  $\lambda' = (\lambda'_1, \lambda'_2, \lambda'_3, \lambda'_4)$ , then*

$$\lambda = A_{m,p} \lambda',$$

*where  $m$  and  $p$  are nonnegative integers and  $A_{m,p}$  is the matrix*

$$\begin{pmatrix} 0 & 0 & 1 & 1 \\ m-1 & m & 0 & 0 \\ p & p & p-1 & p \\ 1 & 1 & 1 & 1 \end{pmatrix}.$$

#### Proof

We make the induction castle of  $X_4 = [P_0, P_4[$ .

Let  $P_2 = T^{-1}\gamma_1$ ; by  $T$ ,  $[P_0, P_2[$  goes to  $[0, \gamma_1[ = X_1$ , which goes by  $T$  to  $[\beta_2, \beta_3[ \subset X_3$ .

On the right of the picture,  $T[P_2, P_4[ = [\gamma_1, \beta_1[ \subset X_2$ . Let  $P_3 = T^{-m}\gamma_2$ , thus  $m$  is the smallest  $k$  such that  $T^{-k}\gamma_2$  is in  $]P_2, P_4[$ . For  $1 \leq k < m$   $T^k[P_2, P_4[ \subset X_2$ , then  $T^m[P_2, P_3[ = [T^m P_2, \gamma_2[ \subset X_2$ , which is further sent by  $T$  to  $[T^{m+1}P_2, \beta_2[ \subset X_3$ , while  $T^m[P_3, P_4[ = [\gamma_2, T^m P_4[ \subset X_3$ . Note that  $T^{m+1}P_2 = T^m P_4$  as the image of  $\gamma_1$  to the right is  $\beta_1$ .

Thus  $J = [\gamma_2, \gamma_2 + \lambda_4[$  is the union of the successive intervals  $T^m[P_3, P_4[$ ,  $T^{m+1}[P_2, P_3[$ ,  $T^2[P_0, P_2[$ . And  $P_1 = T^{-p-2}\gamma_3$  is thus in  $]P_0, P_2[$ . Then  $p$  is the smallest  $k$  such that  $T^{-k}\gamma_3$  is in  $] \gamma_2, \gamma_2 + \lambda_4[$ ,  $J, \dots, T^{p-1}J$  are in  $X_3$  while  $T^p[\gamma_2, T^2 P_1[ \subset X_3$ ,  $T^p[T^2 P_1, \gamma_2 + \lambda_4[ \subset X_4$ , and finally  $T^{p+1}[\gamma_2, T^2 P_1[ \subset X_4$ .

Thus we know the induced map  $S$ : on  $[P_0, P_1[$   $S = T^{p+3}$ , on  $[P_1, P_2[$   $S = T^{p+2}$ , on  $[P_2, P_3[$   $S = T^{m+p+1}$ , on  $[P_3, P_4[$   $S = T^{m+p}$ . The order of the image intervals is, from left to right,  $S[P_1, P_2[$ ,  $S[P_3, P_4[$ ,  $S[P_2, P_3[$ ,  $S[P_0, P_1[$ .

Let  $\lambda'_{4-i}$  be the length of the interval  $[P_i, P_{i+1}[$ ; then we get the required matrix equality, because, for example, the images  $T^i[P_3, P_4[$ , of length  $\lambda'_1$ , are in  $X_2$   $m - 1$  times then in  $X_3$   $p$  times before returning to  $X_4$ , etc...  $\square$

**Lemma 9.** *Let  $\Omega$  be the open positive cone in  $\mathbb{R}^4$ . Then for any pair of positive integers  $(m, p)$  and any vector  $\lambda \in A_{m,p}\Omega$ , there exists a unique  $\lambda'$  such that  $\lambda = A_{m,p}\lambda'$ , and  $T_{\lambda,\pi}$  satisfies all the requirements of Lemma 8 with these values of  $m$  and  $p$ .*

**Proof**

We check that  $A_{m,p}$  is of determinant one and maps  $\Omega$  into  $\Omega$ , and the matrix equality implies the conditions on the lengths of the lemma.  $\square$

**Lemma 10.** *For every infinite sequence of matrices  $A_{m_k,p_k}$ , the set  $\bigcap_{k \in \mathbb{N}} A_{m_1,p_1} \dots A_{m_k,p_k} \Omega$  is nonempty.*

**Proof**

We check that any product of two successive  $A$  has all its entries positive. Let  $\bar{\Omega}$  be the closed positive cone in  $\mathbb{R}^4$ ,  $K_n = A_{m_1,p_1} \dots A_{m_n,p_n} \Omega$ ,  $\bar{K}_n = A_{m_1,p_1} \dots A_{m_n,p_n} \bar{\Omega}$ ,  $K'_n = \bar{K}_n \setminus \{0\}$ ; we have  $K_n \subset K'_n \subset \bar{K}_n$ . But if  $v$  is in  $\bar{\Omega}$  with at least one strictly positive coordinate, then  $A_{m_{n-1},p_{n-1}} \dots A_{m_n,p_n} v$  is in  $\Omega$ , thus

$$\bigcap_{n \geq 1} K_n = \bigcap_{n \geq 1} \bar{K}_n \setminus \{0\} = \bigcap_{n \geq 1} K'_n.$$

Also, each  $K'_n$  is invariant by  $v \rightarrow \lambda v$  for any scalar  $\lambda$ , thus the  $K'_n$  are decreasing compact sets in a projective space, thus their infinite intersection is non-empty; thus  $\bigcap_{n \geq 1} K_n$  is non-empty.  $\square$

**Proposition 11.** *Let  $E$  be the set  $\bigcap_{k \in \mathbb{N}} A_{m_1,p_1} \dots A_{m_k,p_k} \Omega$  normalized by  $\lambda_1 + \dots + \lambda_4 = 1$ . For every  $\lambda \in E$ , there exists a decreasing sequence of intervals  $J_k$  such that the induced map  $S_k$  of the four-interval exchange  $T_{\lambda,\pi}$  on  $J_k$  is the four-interval exchange  $T_{\lambda'_{(k)},\pi}$  (after renormalization, and with reversed order if  $k$  is odd), with  $\lambda = A_{m_1,p_1} \dots A_{m_k,p_k} \lambda'_{(k)}$ .*

*$T_{\lambda,\pi}$  is minimal if the first coordinate of  $\lambda'_{(2k+1)}$  or the last coordinate of  $\lambda'_{(2k)}$  tend to 0 when  $k \rightarrow +\infty$ .*

**Proof**

When we renormalize and reverse the order of the intervals the induced map  $S$  of Lemma 8 is exactly  $T_{\lambda',\pi}$ , and we iterate the construction. At each stage, the induction castle (for  $T$ ), of  $J_k$  fills all the space, thus if it is made of small intervals we can make the reasoning of Proposition 2 to prove minimality.  $\square$

**Proposition 12.** *For any  $\lambda$  in  $E$ , the natural codings of  $T_{\lambda,\pi}$  are adic words. Namely there exist finite words  $B_{n,1}, \dots, B_{n,4}$  such that, for all  $n$ , all the trajectories are infinite concatenations of  $B_{n,i}$ , with*

- $B_{0,i} = i$ ,  $1 \leq i \leq 4$ ,

- for each  $1 \leq i \leq 4$ , there exist an integer  $t(n, i) > 0$ , and  $t(n, i)$  integers  $1 \leq k_s(n, i) \leq 4$  such that

$$B_{n,i} = \prod_{s=1}^{t(n,i)} B_{n-1, k_s(n,i)}$$

The matrix  $A_{m_n, p_n}$  has on its  $i$ -th line,  $1 \leq i \leq 4$ , and  $j$ -th column,  $1 \leq j \leq 4$ , the number of  $1 \leq s \leq t(n, j)$  such that  $k_s(n, j) = i$ .

Each point  $\mu$  in the set  $\bigcap_{k \in \mathbf{N}} A_{m_1, p_1} \dots A_{m_k, p_k} \Omega$  normalized by  $\mu_1 + \dots + \mu_4 = 1$ , defines an invariant probability measure on  $(X, T)$  such that  $\mu(X_i) = \mu_i$ ; every invariant probability measure on  $(X, T)$  is of that form.

### Proof

For  $1 \leq i \leq 4$  we define  $F_{n,i}$  to be the sub-interval of  $J_n$  labelled  $i$  in the definition of  $S_n$ . By construction, for each given  $n$ , the  $T^j F_{n,i}$ ,  $1 \leq i \leq 4$ ,  $0 \leq j \leq |h_{n,i}| - 1$  form a partition of  $X$  into  $k$  Rokhlin towers; these partitions are increasing (the atoms of the  $n + 1$ -th partition are subsets of atoms of the  $n$ -th partition), and, except possibly for a countable number of points, two points belonging to the same atom of the  $n$ -th partition for every  $n$  are the same; we say that the system  $(X, T)$  is of rank at most 4. Moreover, if  $x \in F_{n,4}$ , resp.  $F_{n,3}$ , resp.  $F_{n,2}$ , resp.  $F_{n,1}$ , the nonnegative trajectory of  $x$  under  $S_{n-1}$ , for the natural coding given by Proposition 11, begins with  $413^{p_n+1}$ , resp.  $413^{p_n}$ , resp.  $42^{m_n} 3^{p_n}$ , resp.  $42^{m_n} 3^{p_n-1}$ . By iterating this process, if  $x \in F_{n,i}$ , the nonnegative trajectory of  $x$  under  $T$  begins with a word denoted by  $B_{n,i}$ . It follows from the definitions that  $F_{n-1,i}$  is a union of images by  $T$  of the  $F_{n,j}$ ,  $1 \leq j \leq k$ , and thus the  $B_{n,i}$  are made by the above concatenation rules, and we check the matrix.

This is enough to ensure that a measure on  $X_u$  is determined by its values on the atoms of these partitions, thus, if it is  $T$ -invariant, by its values on the  $F_{n,i}$ . We check also that if  $v_n = (\mu(F_{n,1}), \dots, \mu(F_{n,k}))$ , we get  $v_{n-1} = A_{m_n, p_n} v_n$ . Thus the measure  $\mu$  is completely determined by the vector  $v_0$ .  $\square$

**Proposition 13.** *If  $p_1 \geq 9$  and  $3(p_n + 1) \leq m_n \leq \frac{1}{2}(p_{n+1} + 1)$  for all  $n$ ,  $T$  is minimal and not uniquely ergodic.*

### Proof

The condition of minimality is satisfied as  $p_n \rightarrow +\infty$  and every coordinate of  $\lambda'_n$  is smaller than  $\frac{1}{p_n-1}$ .

We define  $M_n = A_{m_n, p_n}$ ,  $\tilde{M}_n x = \frac{M_n x}{|M_n x|}$  where  $|y| = \sum_{i=1}^4 y_i$ . Let  $\tilde{B}_k$  be the mapping  $\tilde{M}_1 \dots \tilde{M}_k$ .

We note that for  $i = 1$  or  $i = 4$ , and any  $x$ ,  $(M_n x)_i \leq 1$  while  $|M_n(x)| \geq p_n + 1$  thus  $(\tilde{M}_n x)_i \leq \frac{1}{p_n+1}$ .

Suppose  $2m_n \leq p_{n+1} + 1$  for all  $n$  and let  $e_3 = (0, 0, 1, 0)$ ; we prove that then for  $n \geq 1$ ,  $(\tilde{B}_n e_3)_3 \geq 1 - \frac{3}{p_1+1}$ . Let  $x^{k+1} = e_3$ ,  $x^{j-1} = \tilde{M}_{j-1} x^j$ , for  $2 \leq j \leq k+1$ ; then, in view of the previous result, it is enough to prove that  $x_2^j \leq \frac{1}{p_1+1}$  for  $j = 1$ , and we shall prove it by induction on  $j$ ; this is true for  $j = k+1$ , and under the induction hypothesis

$$x_2^j = \frac{(m_j - 1)x_1^{j+1} + m_j x_2^{j+1}}{p_j + 1 + (m_j - 1)x_1^{j+1} + m_j x_2^{j+1} + x_4^{j+1}} \leq \frac{2m_j}{(p_j + 1)(p_{j+1} + 1)} \leq \frac{1}{p_j + 1}.$$

Suppose  $m_n \geq 3(p_n + 1)$  and let  $e_2 = (0, 1, 0, 0)$ ; we prove that for  $n \geq 1$ ,  $(\tilde{B}_n e_2)_2 \geq \frac{1}{3}$ . We define a sequence  $x_i$  in the same way as in the previous paragraph, with the induction hypothesis  $x_2^{j+1} \geq \frac{1}{3}$ ,

We define now  $\mu$  by a vector which is a limit (on a subsequence) of the  $A_{m_1, p_1} \dots A_{m_k, p_k} e_2$ , and  $\nu$  in the same way with  $A_{m_1, p_1} \dots A_{m_k, p_k} e_3$ . Then if  $\mu = \nu$ , it would give measure at least  $1 - \frac{3}{n_1+1}$  to the set  $X_3$  and at least  $\frac{1}{3}$  to the set  $X_2$  which is a contradiction as soon as  $n_1 \geq 9$ .  $\square$

These examples can satisfy the requirement of total irrationality, which was not satisfied by Veech's examples: it is proved in [17] that for any given hyperplane  $H$  we can find sequences  $(m_n, p_n)$  satisfying the conditions of Proposition 13 and such that  $\bigcap_{k \in \mathbb{N}} A_{m_1, p_1} \dots A_{m_k, p_k} \Omega$  does not intersect  $H$ , and we can avoid a countable family of hyperplanes.

There is a duality between the lengths of the intervals and the values of the invariant measures on them: with the notations of Section 4, for any  $\lambda \in E$ , every invariant probability measure on  $([0, 1[, T_{\lambda, \pi})$  is defined from a vector  $\mu \in E$  by giving measure  $\mu_i$  to the  $i$ -th interval. Under the above conditions on the  $m_k, p_k$ ,  $E$  is not reduced to a point but is a segment, whose two endpoints give the two invariant ergodic measures. If we choose  $\lambda$  to be in the interior of this segment, these two ergodic measures are absolutely continuous with respect to the Lebesgue measure but different from it; if we choose  $\lambda$  to be an endpoint, one ergodic measure is the Lebesgue measure and the other one is singular; a recent work of J. Chaika [6] has proved that this singular measure can have a support of arbitrarily small Hausdorff dimension.

## 5. UNIQUE ERGODICITY AFTER BOSHERNITZAN

We call  $m$  the normalized Lebesgue measure on  $\Lambda_r$ , and  $\rho$  the Lebesgue measure on  $[0, 1[$ .

**Theorem 14.** *For a given irreducible  $\pi$ ,  $T_{\lambda, \pi}$  is uniquely ergodic for  $m$ -almost every  $\lambda \in \Lambda_r$ .*

This result was the first big conjecture on iet, stated as a question in [17] and proved independently by W. Veech [25] and H. Masur [18]. The proofs of Veech and Masur use elaborate geometric methods; but a later proof of M. Boshernitzan uses mainly combinatorial methods; it is published in [2] but can be simplified (and made purely combinatorial) by using [4]. Thus we give here this simplified proof, with an updated vocabulary: in particular, the Rauzy graphs are used without being named im [2], and as far as we know this is the first published paper where they are mentioned.

**Definition 10.** *Let  $(X_L, S)$  be a minimal symbolic system. If  $\mu$  is an  $S$ -invariant probability measure, for each natural integer  $n$ , we denote by  $e_n(S, \mu)$  the smallest positive measure of the cylinders defined by words of length  $n$  of  $L$ .*

**Proposition 15.** [4] *If for some invariant probability measure  $\mu$ ,  $ne_n(\mu)$  does not tend to 0 when  $n$  tends to  $+\infty$ , then the system  $(X_L, S)$  is uniquely ergodic.*

### Proof

Then for infinitely many  $n$  we have  $e_n \geq \frac{a}{n}$  and thus  $p(n) \leq (C-1)n$ , thus  $\liminf_{n \rightarrow +\infty} p(n) - Cn = -\infty$ . Thus on a subsequence  $p(n+1) - C(n+1) \leq p(n) - Cn$  and  $p(n+1) - C(n+1) \leq 0$ . On this subsequence we have both  $p(n) \leq p(n+1) \leq C(n+1) \leq C'n$  for  $n$  large enough and  $p(n+1) - p(n) \leq C$  thus at most  $C$  right special words. By the reasoning of the first part of the proof of Proposition 6 we conclude that there is a finite number of ergodic invariant measures.

Thus if  $(X_L, S)$  is not uniquely ergodic there are two invariant ergodic measures  $\mu_i$  and a finite word  $w$  such that  $\mu_1[w] < \mu_2[w]$  and in the interval  $]\mu_1[w], \mu_2[w][$  there is no  $\nu[w]$  for  $\nu$  any ergodic invariant measure. We fix  $\mu_1[w] < t < s < \mu_2[w]$ .

Let  $E_n = \{x; \frac{N(w, x_0 \dots x_{n-1})}{n} \in ]t, s[ \}$ . As  $\frac{N(w, x_0 \dots x_{n-1})}{n} \rightarrow \nu[w]$   $\nu$ -almost everywhere, we get  $\nu(E_n \text{ infinitely often}) = 0$ , thus  $\nu(E_n) \rightarrow 0$  when  $n \rightarrow +\infty$ , thus also  $\mu(E_n) \rightarrow 0$  by convex combination.

By genericity, we choose  $x$  and  $y$  such that  $\frac{N(w, x_0 \dots x_{n-1})}{n} < t < s < \frac{N(w, y_0 \dots y_{n-1})}{n}$  for  $n$  large enough. We choose in the Rauzy graph of length  $n$  a path from  $x_0 \dots x_{n-1} = v_1$  to  $y_0 \dots y_{n-1} = v_p$ , through  $v_i$ ,  $2 \leq i \leq p-1$ ; it exists by minimality and can be chosen of minimal length thus without loops.

Then  $|N(w, v_i) - N(w, v_{i+1})| \leq 1$ . To go from below  $tn$  to above  $sn$  by moving by 1 at a time, we need to be at least  $n(s-t) - 2$  times between  $tn$  and  $sn$ . Thus  $[v_i]$  is in  $E_n$  for at least  $n(s-t) - 2$  values of  $i$ , and the  $v_i$  are all distinct. We get  $\mu(E_n) \geq (n(s-t) - 2)e_n$  and thus  $ne_n \rightarrow 0$ , contradiction.  $\square$

For a given iet  $T_{\lambda, \pi}$ , we call  $\rho$  the Lebesgue measure on the interval  $[0, 1[$ , and call  $e_n(T_{\lambda, \pi})$  the quantity  $e_n(S, \rho)$  defined for the associated symbolic system.

**Proposition 16.** [2] *Let  $U_{n, \epsilon}$  be the set of  $\lambda \in \Lambda_r$  such that  $e_n(T_{\lambda, \pi}) \leq \frac{\epsilon}{n}$ . If  $\epsilon$  is small enough,  $m(U_{n, \epsilon}) \leq 3r^3\epsilon$ .*

### Proof

Let  $G_n(\lambda)$  be the Rauzy graphs of length  $n$  of the language of  $T_{\lambda, \pi}$ . As the complexity of any  $u(x)$  is  $(r-1)n + 1$ , by Lemma 4  $G_n$  has at most  $3r - 3$  branches, as a branch starts at a left or right special factor, and there are at most respectively  $r - 1$  and  $2r - 2$  in each case.

We recall that  $\psi$  is a *weight function* on a graph if it is positive on each vertex, the sum of its values on vertices in 1, and it can be extended to the edges such that for every vertex

$$\psi(w) = \sum_{\text{incoming edges}} \psi(e) = \sum_{\text{outcoming edges}} \psi(e).$$

We define a function  $\psi_\lambda$  on the vertices of  $G_n(\lambda)$  by associating to the vertex  $w_1 \dots w_n$  the measure of the cylinder,  $\rho[w_1 \dots w_n]$ ;  $\psi_\lambda$  is a weight function on the graph  $G_n(\lambda)$ , and the weight of an edge  $w_1 \dots w_{n+1}$  is also  $\rho[w_1 \dots w_{n+1}]$ .

We fix now a Rauzy graph  $G$  of length  $n$ ; let  $\Lambda(G)$  be the set of  $\lambda \in \Lambda_r$  such that  $G_n(\lambda) = G$ . For a given word  $w = w_1 \dots w_n$ ,  $\psi_\lambda(w_1)$  is just  $\lambda_{w_1}$ ; for all  $\lambda \in \Lambda(G)$ , all the Rauzy graphs  $G_i(\lambda)$ ,  $1 \leq i \leq n$ , are fixed, and determine the way the measures of cylinders of length  $i+1$  are computed from those of length  $i$ , through the defining equalities of the weight function  $\psi_\lambda$  on  $G_i(\lambda)$ ; thus the numbers  $\psi_\lambda(w_1 \dots w_i)$ ,  $1 < i \leq n$  can be computed inductively; they depend linearly on  $\lambda$ . Because  $T_{\lambda, \pi}$  preserves the measure  $\rho$ ,  $\psi_\lambda(w_1 \dots w_n) = \psi_\lambda(w'_1 \dots w'_n)$  if  $w_1 \dots w_n$  and  $w'_1 \dots w'_n$  are on the same branch of  $G$ ; hence, for fixed  $\lambda$ ,  $\psi_\lambda(w_1 \dots w_n)$  takes  $1 \leq t \leq 3r - 3$  values, which we denote by  $\phi_1(\lambda), \dots, \phi_t(\lambda)$ ; the  $\phi_j$  are linear functionals,  $e_n(T_{\lambda, \pi})$  is just the smallest of the  $\phi_j(\lambda)$ ,  $1 \leq j \leq t$ . Furthermore, again through the defining equalities of the successive weight functions on  $G_i(\lambda)$ , we can retrieve  $\lambda$  from the values  $\psi_\lambda(w)$  on all the vertices of  $G$ ; thus  $\Lambda(G)$  is a convex set and every weight function  $\psi$  on  $G$  yields a  $\lambda \in \Lambda(G)$  such that  $\psi_\lambda = \psi$ .

We want to estimate the measures of  $\{\lambda \in \Lambda(G); \phi_i(\lambda) \leq \frac{\epsilon}{n}\}$ ; for this, we use a general result for which we refer the reader to [3], Corollary 7.4: *If  $\phi$  is the restriction of a linear functional to a*

convex set  $K$  of dimension  $d$ , taking values between 0 and  $A$ , then, if  $V$  denotes the volume,

$$V(\pi^{-1}[0, B]) \leq \frac{dB}{A}V(K).$$

We apply it with  $K = \Lambda(G)$ , restricting ourselves to those with  $m(\Lambda(G)) > 0$ ,  $\phi = \phi_i$ ,  $B = \frac{\epsilon}{n}$ ; the dimension is  $r - 1$ , the volume is the Lebesgue measure; we need an estimate on  $A$ ; for this, we claim that *for each vertex  $s$  of  $G$ , there exists a weight function such that  $\psi(s) \geq \frac{1}{rn}$* . To do this, we choose a  $\lambda \in \Lambda(G)$  such that  $T_{\lambda, \pi}$  is minimal, which is possible as  $m(\Lambda(G)) > 0$ ; this implies that  $G$  is strongly connected and thus we can find a loop  $s_0 \rightarrow \dots s_k \rightarrow s_0$  in  $G$ ; by taking it of minimal length, we ensure it has no repetition. Then we define  $\psi'$  to be  $\frac{1}{k+1}$  on the  $s_i$  and 0 on the other vertices;  $\psi'$  is not a weight function as it may be 0 on some vertices, but  $\psi = (1 - \delta)\psi' + \delta\psi_\lambda$  is a weight function, and as  $k \leq (r - 1)n + 1$  we can choose  $\delta$  such that our claim is proved.

Thus we have  $A \geq \frac{1}{rn}$ , and thus, for all  $G$  with  $m(\Lambda(G)) > 0$  and hence for all  $G$ ,

$$m(\{\lambda \in \Lambda(G); \phi_i(\lambda) \leq \frac{\epsilon}{n}\}) \leq (r - 1)r\epsilon m(\Lambda(G)).$$

As  $t \leq 3r - 3$ ,

$$m(\{\lambda \in \Lambda(G); \min_{1 \leq i \leq t} \phi_i(\lambda) \leq \frac{\epsilon}{n}\}) \leq 3(r - 1)^2 r \epsilon m(\Lambda(G)),$$

which implies the proposition.  $\square$

#### Proof of Theorem 14

For small  $0 < \epsilon$  and  $n \geq 1$ , we put  $V_{n, \epsilon} = \Lambda_r \setminus U_{n, \epsilon}$ , and  $V_\epsilon = \bigcap_{N \geq 1} \bigcup_{n > N} V_{n, \epsilon} \cap \{\lambda; T_{\lambda, \pi} \text{ i.d.o.c.}\}$ .

If  $\lambda$  is in  $V_\epsilon$ , there are infinitely many  $n$  such that  $e_n(T_{\lambda, \pi}, \rho) \geq \frac{\epsilon}{n}$ , hence  $ne_n(T_{\lambda, \pi}, \rho) \not\rightarrow 0$  when  $n \rightarrow +\infty$ , and  $T_{\lambda, \pi}$  is uniquely ergodic by Proposition 15. Thus  $m(\{\lambda; T_{\lambda, \pi} \text{ is uniquely ergodic}\})$  is at least  $m(V_\epsilon) \geq 1 - 3r^3\epsilon$ , and thus is one as  $\epsilon$  is arbitrary.  $\square$

The above proof does not use any of the geometric properties of interval exchange maps; what it needs is only that it is a class of symbolic systems on an  $r$ -letter alphabet, of complexity at most  $sn$  for a fixed  $s$ , with a common invariant measure  $\rho$  such that the vector  $(\rho[1], \dots, \rho[r])$  takes all possible values in  $\Lambda_r$ .

## 6. WEAK MIXING AFTER BOSHERNITZAN [5]

The second big conjecture on iet was about weak mixing: for a given  $\pi$  outside the so-called rotation class,  $T_{\lambda, \pi}$  is weakly mixing for  $m$ -almost every  $\lambda \in \Lambda_r$ . This resisted more than twenty years before being proved by A. Avila and G. Forni [1]. The recent result we give now is not known to be equivalent, but has at least a similar flavour and its proof is quite short.

**Definition 11.**  $\phi(x)$  is the largest  $s$  such that for infinitely many  $n$  there is a Rokhlin tower of total measure  $4s$ , made of  $2n + 1$  intervals, with  $x$  in the middle of the middle level.

We check that if  $D$  be the set of  $\gamma_i$ , for  $1 \leq i \leq r - 1$  such that  $\pi(i + 1) \neq \pi i$ , we have

$$\rho_n(x) = \frac{1}{2} \min_{|p \leq n, |q| \leq n, p \neq q} |T^p x - T^q x| \wedge \min_{-n \leq k \leq n} \min_{\gamma_i \in D} |T^k x - \gamma_i|,$$

$$\phi(x) = \limsup_{n \rightarrow +\infty} n \rho_n(x).$$

**Proposition 17.** *If  $T$  is ergodic for the Lebesgue measure  $\rho$ , if  $\phi(t) > 0$ , the induced map of  $T$  on  $[0, t[$  is weakly mixing for the Lebesgue measure on  $[0, t[$ .*

**Proof**

For infinitely many  $m$ , we build a tower of total measure  $> a > 0$ , made of  $2m + 1$  intervals, with  $t$  in the middle of the middle level. The intersection of this tower with  $[0, t[$  consists of about  $mt$  (by the ergodic theorem for  $T^{-1}$ ) full levels below the one containing  $t$ , the left half of the level containing  $t$ , and about  $mt$  (by the ergodic theorem for  $T$ ) full levels above the one containing  $t$ ; we trim it to get exactly  $n$  levels, called  $Y_{n,k}$ , above and below.

Suppose the induced map  $S$  has an eigenfunction  $f$  for the eigenvalue  $\theta$ . Because the  $Y_{n,k}$  are small intervals, there exists a sequence of maps  $f_n$  such that  $\|f - f_n\|_1 \rightarrow 0$  and  $f_n$  has a constant value  $f_{n,k}$  on each  $Y_{n,k}$ . The  $f_{n,k}$  can be taken of modulus 1.

Let  $Z$  be the tower made with the left halves of the  $Y_{n,k}$ ,  $-n + 1 \leq k \leq -1$ ; it has total measure at least  $\frac{a}{5}$ . If  $r_n$  is the translation taking the left half to the right half of each  $Y_{n,k}$ , and if  $x$  is in a level of  $Z$ , we have  $f_n(S^{n+1}x) = f_n(S^n r_n x)$ . Then, for any given  $\epsilon$  and  $n$  large enough,

$$\begin{aligned} \rho(Z)|\theta - 1| &= \sum_{k=-n+1}^{-1} |\theta^{n+1} f_{n,k} - \theta^n f_{n,k}| \rho(Y_{n,k}) = \int_Z |\theta^{n+1} f_n(x) - \theta^n f_n(r_n x)| dx \leq \\ &2\epsilon + \int_Z |\theta^{n+1} f(x) - \theta^n f(r_n x)| dx = 2\epsilon + \int_Z |f(S^{n+1}x) - f(S^n r_n x)| dx \leq \\ &4\epsilon + \int_Z |f_n(S^{n+1}x) - f_n(S^n r_n x)| dx = 4\epsilon. \end{aligned}$$

Thus  $\theta = 1$ , which is excluded as  $S$  is ergodic. □

**Proposition 18.** *If  $T$  is minimal and ergodic for the Lebesgue measure, the set of  $t$  such that  $\phi(t) > 0$  is residual and of full Lebesgue measure.*

**Proof**

By making the induction castle of a small subinterval, for any given  $N$  we can make towers of at least  $N$  intervals of total measure at least  $\frac{1}{r+2}$ . Let  $Y_n$  be a sequence of such towers, with  $h_n$  intervals, and  $Z_n$  be the middle ninth (=middle third in length and height) of  $Y_n$ . Let  $Z$  be  $\{Z_n \text{ infinitely often}\}$ .  $Z$  is residual and  $\rho(Z)$  is at least  $\frac{1}{10(r+2)}$ , while if  $z$  is in  $Z_n$  there is a Rokhlin tower of total measure at least  $\frac{1}{6(r+2)}$ , made of  $h_n \rightarrow +\infty$  intervals, with  $z$  in the middle of the middle level.

As the set of  $t$  such that  $\phi(t) > 0$  is  $T$ -invariant, we conclude by ergodicity. □

## 7. SIMPLICITY: THE LAST FRONTIER?

The third big conjecture on iet is still open, and we end this course by stating it as Question 1 below, with the necessary historical background.

One recurrent preoccupation of ergodicists in the last twenty years has been with joinings: the notion of *self-joinings* of a system has been introduced by D. Rudolph in [21], to generalize some useful invariants of measure-theoretic isomorphism such as the factor algebra and the centralizer.

**Definition 12.** *A self-joining (of order two) of a system  $(X, T, \mu)$  is any measure  $\nu$  on  $X \times X$ , invariant under  $T \times T$ , for which both marginals are  $\mu$ .*

**Definition 13.** An ergodic system  $(X, T, \mu)$  has minimal self-joinings (of order two) if any ergodic self-joining (of order two)  $\nu$  is either the product measure  $\mu \times \mu$  or a diagonal measure defined by  $\nu(A \times B) = \mu(A \cap T^i B)$  for an integer  $i$ .

A transformation which has minimal self-joinings has trivial centralizer and no nontrivial proper factor, and can be used to build a so-called *counter-example machine* with surprising properties. The first example of a transformation with minimal self-joinings was given in [21], and a little later the famous *Chacon map* was shown in [8] also to have minimal self-joinings. However, both these examples may seem built on purpose, and they have no “natural”, i.e. geometric, realization. Then geometric examples of transformations with minimal self-joinings were sought in the category of interval exchange transformations. And indeed in 1983 A. del Junco [7] built a one-parameter family of three-interval exchange transformations, depending on an irrational  $\gamma$ , and proved that whenever this  $\gamma$  has bounded partial quotients in its continued fraction expansion the system has minimal self-joinings (the interested reader is warned that he will *not* find the terms “three-interval exchange transformation” or “minimal self-joinings” in del Junco’s paper; the systems which he describes as two-point extensions of rotations are indeed three-interval exchange transformations, and the notion of “simplicity” he proves is only slightly weaker than the original notion of minimal self-joinings, and has been standing as the current definition of “minimal self-joinings” since [9]).

But in the meantime, Veech [26] had shown that almost all interval exchange transformation (in the sense: for a fixed permutation, for Lebesgue-almost all values of the lengths of the intervals) is rigid:

**Definition 14.** A system  $(X, T, \mu)$  is rigid if there exists a sequence  $s_n \rightarrow \infty$  such that for any measurable set  $A$

$$\mu(T^{s_n} A \Delta A) \rightarrow 0.$$

Simple systems have uncountable centralizers and cannot have minimal self-joinings; thus Veech devised in [24] a weakened notion of minimal self-joinings to allow for a nontrivial centralizer; the new notion, which Veech called “property S”, is now known as “simplicity (in the sense of Veech)”:

**Definition 15.** An ergodic system  $(X, T, \mu)$  is simple of order two if any ergodic self-joining of order two  $\nu$  is either the product measure  $\mu \times \mu$  or a measure defined by  $\nu(A \times B) = \mu(A \cap S^{-1} B)$  for some measurable transformation  $S$  commuting with  $T$ .

Simplicity is strong enough to keep many of the properties of systems with minimal self-joinings, though proving this required a lot of work [10] [24]. And Veech asked the following question (4.9 of [24])

**Question 1.** Are almost all interval exchange transformations simple?

The notion of simplicity having been devised just for that, it is tacitly conjectured that indeed almost all interval exchange transformations are weakly mixing, simple and rigid.

But, while Veech’s question stood unanswered, examples of simple transformations remained scarce: there were of course the systems with minimal self-joinings, and some systems without minimal self-joinings but naturally related to these systems (such as the time-one map of a flow which, as a flow, has minimal self-joinings); at last in [9], a natural generalization of Chacon’s map was (very cunningly!) shown to be simple and rigid. It remains to this day the main explicit example of a simple and rigid map; among interval exchange transformations, some more 3-iet, beside del Junco’s, were proved to have minimal self-joinings [11]; and at long last some 3- [11] and 4- [12] interval exchanges were proved to be simple and rigid, but they are so because they are

measure-theoretically isomorphic to del Junco-Rudolph' map.

Thus an answer to Veech's question seems still to be a very difficult problem, which also fell a little out of fashion; the result of Avila-Forni was a necessary step towards a positive answer, but their proof of weak mixing does not seem to imply anything in the direction of simplicity. However, the result of Boshernitzan in the previous section of the present course may give a faint glimmer of new hope, as it proves weak mixing by the "Chacon trick" of building two towers of heights differing by one, and it is this trick which implies the weak mixing of Chacon's and del Junco-Rudolph' maps, but also the minimal self-joinings of Chacon's map and (after long manipulations using the full knowledge of the system, and not only local towers), the simplicity of del Junco-Rudolph' map.

#### REFERENCES

- [1] A. AVILA, G. FORNI: Weak mixing for interval exchange maps and translation flows, *Ann. of Math. (2)* 165 (2007), p. 637–664.
- [2] M. BOSHERNITZAN: A condition for minimal interval exchange maps to be uniquely ergodic, *Duke Math. J.* 52 (1985), p. 723–752.
- [3] M. BOSHERNITZAN: A unique ergodicity of minimal symbolic flows with linear block growth, *J. Analyse Math.* 44 (1984/85), p. 77–96.
- [4] M. BOSHERNITZAN: A condition for unique ergodicity of minimal symbolic flows, *Ergodic Theory Dynam. Systems* 12 (1992), p. 425–428.
- [5] M. BOSHERNITZAN: A criterion for weak mixing of induced interval exchange transformations, arXiv:1105.0239v1.
- [6] J. CHAIKA: Hausdorff dimension for ergodic measures of interval exchange transformations. *J. Mod. Dyn.* 2 (2008), p. 457–464.
- [7] A. del JUNCO: A family of counterexamples in ergodic theory, *Israel J. Math.* 44 (1983), p. 160–188.
- [8] A. del JUNCO, A.M. RAHE, M. SWANSON: Chacon's automorphism has minimal self-joinings, *J. Analyse Math.* 37 (1980), p. 276–284.
- [9] A. del JUNCO, D.J. RUDOLPH: A rank-one, rigid, simple, prime map, *Ergodic Th. Dyn. Syst.* 7, 2 (1987), p. 229–247.
- [10] A. del JUNCO, D.J. RUDOLPH: On ergodic actions whose self-joinings are graphs, *Ergodic Th. Dyn. Syst.* 7, 4 (1987), p. 531–557.
- [11] S. FERENCZI, C. HOLTON, L. ZAMBONI: Joinings of three-interval exchange transformations, *Ergodic Th. Dyn. Syst.* 25 (2005), p. 483–502.
- [12] S. FERENCZI, L.Q. ZAMBONI, Eigenvalues and simplicity of interval exchange transformations, *Ann. Sci. Ec. Norm. Sup. (4)* 44 (2011), p. 361–392.
- [13] A.B. KATOK: Invariant measures of flows on orientable surfaces, *Dokl. Akad. Nauk. SSSR* 211 (1973), p. 775–778 (in Russian).
- [14] A. KATOK: Interval exchange transformations and some special flows are not mixing, *Israel J. Math.* 35 (1980), p. 301–310.
- [15] A.B. KATOK, E.A. SATAEV: Standardness of automorphisms of transposition of intervals and fluxes on surfaces, *Math. Zametki* 20, 4 (1976), p. 479–488 (in Russian), translated in *Math. Notes Acad. Sci. USSR* (1977), p. 826–831.
- [16] M.S. KEANE: Interval exchange transformations, *Math. Zeitsch.* 141 (1975), p. 25–31.
- [17] M.S. KEANE: Non-ergodic interval exchange transformations, *Israel J. Math.* 26 (1977), p. 188–196.
- [18] H. MASUR: Interval exchange transformations and measured foliations, *Ann. of Math.* 115 (1982), p. 169–200.
- [19] M. MORSE, G. A. HEDLUND: Symbolic Dynamics, *Amer. J. Math.* 60 (1938), p. 815–866.
- [20] V.I. OSELEDETS: On the spectrum of ergodic automorphisms, *Doklady Akademii Nauk. SSSR* 168, 5 (1966), p. 1009–1011 (in Russian), translated in *Soviet Math. Doklady* 7 (1966), p. 776–779.
- [21] D.J. RUDOLPH: An example of a measure-preserving map with minimal self-joinings, and applications, *J. Analyse Math.* 35 (1979), p. 97–122.

- [22] W.A. VEECH: Strict ergodicity in zero dimensional dynamical systems and the Kronecker-Weyl theorem mod. 2, *Trans. Amer. Math. Soc.* 140 1969, p. 1–33.
- [23] W.A. VEECH: Interval exchange transformations, *J. Analyse Math.* 33 (1978), p. 222–272.
- [24] W.A. VEECH: A criterion for a process to be prime, *Monatsh. Math.* **94**, 4 (1982), p. 335–341.
- [25] W.A. VEECH: Gauss measures for transformations on the space of interval exchange maps, *Ann. of Math.* 115 (1982), p. 201–242.
- [26] W.A. VEECH: The metric theory of interval exchange transformations I, *Amer. J. Math.* **106** (1984), p. 1331–1358.
- [27] M. VIANA: Dynamics of interval exchange maps and Teichmüller flows, preliminary manuscript available from <http://w3.impa.br/~viana/out/ietf.pdf>.
- [28] J.-C. YOCCOZ: Échanges d’intervalles (in French), *Cours au Collège de France* (2005), [http://www.college-de-france.fr/media/equ\\_dif/UPL8726\\_yoccoz05.pdf](http://www.college-de-france.fr/media/equ_dif/UPL8726_yoccoz05.pdf)

INSTITUT DE MATHÉMATIQUES DE LUMINY, CNRS - UPR 9016, CASE 907, 163 AV. DE LUMINY, F13288 MARSEILLE CEDEX 9 (FRANCE), AND FÉDÉRATION DE RECHERCHE DES UNITÉS DE MATHÉMATIQUES DE MARSEILLE, CNRS - FR 2291

*E-mail address:* `ferenczi@iml.univ-mrs.fr`